

Cola intake increases exposure of erlotinib

10 February 2016



"Potentially, the effects of cola on erlotinib exposure may be extrapolated to other tyrosine kinase inhibitors with a pH-dependent solubility (e.g., dasatinib, gefitinib, nilotinib), but this remains to be evaluated in future studies," the authors write. "Furthermore, other acidic beverages (i.e., orange juice, other carbonated drinks) may have similar effects as cola and should be explored in future trials."

More information: <u>Abstract</u> <u>Full Text (subscription or payment may be required)</u>

Copyright © 2016 HealthDay. All rights reserved.

(HealthDay)—Cola intake leads to a clinically relevant and statistically significant increase in the bioavailability of erlotinib during esomeprazole treatment, according to a study published online Feb. 8 in the *Journal of Clinical Oncology*.

Roelof W.F. van Leeuwen, Pharm.D., from the Erasmus Medical Center Cancer Institute in Rotterdam, Netherlands, and colleagues conducted a randomized, cross-over, pharmacokinetic study of 28 patients with non-small-cell lung cancer. Intrapatient differences were evaluated for absorption (area under the plasma concentration time curve [AUC_{0-12h}]) after a seven-day period of concomitant treatment with erlotinib, with or without esomeprazole, with either cola or water.

The researchers found that in patients treated with erlotinib and esomeprazole with cola, the mean AUC_{0-12h} increased 39 percent (P = 0.004). However, in patients not treated with esomeprazole, the mean AUC_{0-12h} was only slightly higher (9 percent; P = 0.03) after erlotinib intake with cola.



APA citation: Cola intake increases exposure of erlotinib (2016, February 10) retrieved 30 May 2022 from https://medicalxpress.com/news/2016-02-cola-intake-exposure-erlotinib.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.