

Only half of kidney stone patients compliant with Tx

22 February 2016



0.59), and surgery for stone disease (OR, 0.77), compared to nonadherent [patients](#).

"To improve adherence further research is needed to understand patient and provider level factors that contribute to lower rates of adherence," the authors write.

One author disclosed financial ties to the medical device industry.

More information: [Abstract](#)
[Full Text](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

(HealthDay)—Just over half of kidney stone patients are adherent to preventive pharmacological therapy, and adherent patients have lower levels of health care utilization, according to a study published in the March issue of *The Journal of Urology*.

Casey A. Dauw, M.D., from the University of Michigan in Ann Arbor, and colleagues assessed [medication adherence](#) among adult patients with a diagnosis of [kidney stones](#). By accessing medical and pharmacy claims data, [adherence](#) to therapy within the first six months of initiating treatment was measured using the proportion of days covered formula. Adherence was defined as a proportion of days covered value of 80 percent or greater.

The researchers found that just over half (51.1 percent) of 8,950 eligible patients were adherent to preventive pharmacological therapy. Adherent patients had lower odds of emergency department visits (odds ratio [OR], 0.73), hospitalization (OR,

APA citation: Only half of kidney stone patients compliant with Tx (2016, February 22) retrieved 12 October 2022 from <https://medicalxpress.com/news/2016-02-kidney-stone-patients-compliant-tx.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.