

Some drug addicts more likely to relapse than others, study finds

April 13 2016

People with drug addictions who started opioid abuse later in life use injections for their drugs, or increased their use of downers before starting drug treatment, are more likely to relapse from treatment than others, says a new study from McMaster University.

But, the older the person with drug abuse issues, the more likely they will not [relapse](#) from [treatment](#).

"We can improve our tailoring of treatment to each patient if we know who among patients taking methadone treatment is at high risk for opioid relapse," said Dr. Zena Samaan, principal author of the study and an associate professor of psychiatry and behavioural neurosciences of McMaster's Michael G. DeGroote School of Medicine.

"As well, [health care providers](#) can target more aggressive therapies to those at [high risk](#)."

Canada and the U.S. are the world's highest consumers of [prescription opioids](#) which are common drugs for pain management. At least one study says one in six Canadians using prescription opioids. But these medications are also highly addicting and liable for abuse. Methadone maintenance treatment is the most common intervention for those with drug addiction, but relapse is common, with 46% of patients continuing to use illicit opioids during or after the methadone treatment.

"Since opioid disorder is chronic, remitting and relapsing, we wanted to

find those factors that led to longer abstinence from illicit opioids," said Leen Naji, a student of McMaster's Michael G. DeGroot School of Medicine and first author of the paper. "There has been little research on this issue of how long a patient can go without the illicit opioid use."

Samaan and her team looked at 250 adults who had been on a methadone treatment for an average of four years at 13 clinical sites in Ontario.

They found that:

- Those who injected drugs were more than twice as likely to relapse by using opioids while on treatment, than those who did not inject drugs;
- For every year increase in age of starting to abuse opioids there is a 10% increase in relapse;
- For every day of benzodiazepine use in the previous month there is a 7% increase in relapse; while current age carries 7% reduction in risk of relapse (the older the person, the less likely of relapse).

Provided by McMaster University

Citation: Some drug addicts more likely to relapse than others, study finds (2016, April 13)
retrieved 15 May 2023 from <https://medicalxpress.com/news/2016-04-drug-addicts-relapse.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.