


Natural births after caesarean more likely if you call the midwife

20 April 2016, by Jamie Brown



Pregnancy test. Credit: public domain

Women who had a caesarean section in a previous pregnancy are much more likely to have a safer vaginal birth if their antenatal care is led by a midwife, according to a new study from The University of Manchester.

Although caesarean sections are safe, research is increasingly showing that vaginal birth and labour protects against long term-risks such as impaired immune response, asthma, obesity and type 2 diabetes in the baby. As a result there is a growing emphasis placed by health authorities on promoting vaginal birth after caesarean (VBAC) where it is safe to do so.

The new Manchester study, published in the journal *Birth*, is the first to evaluate the influence on VBAC rates of midwife-led antenatal care as opposed to care led by an obstetrician.

Using patient records, 405 women giving birth at one hospital after a previous caesarean were examined in two groups. The first group was from 2008 when antenatal care was led by obstetricians and, following a change in hospital policy in 2009-2010, the second group was drawn from women who had received midwife-led antenatal care in 2011.

The number of women who intended to give birth by VBAC in 2011 was 90.3 percent, against 77 percent in 2008. Those who actually achieved VBAC was also higher in the midwife-led group with 61.2 percent instead of 46.9 percent. Both represented significant increases in VBAC when antenatal care was led by a midwife.

Dr Helen White, a midwifery lecturer in the University's School of Nursing, Midwifery and Social Work, led the research. She said: "There were few differences between the two groups of women we studied, so the main variance was the professional responsible for their care.

"Where it can be achieved safely, vaginal birth is preferable, but there's a real issue with women who have had a caesarean once, opting for the same again. There aren't many initiatives out there to break this cycle so this finding is important for providing evidence that midwives are best placed to promote [vaginal birth](#)."

The authors of the study are not proposing that women should reject [caesarean birth](#) when they are advised to do so, but suggest that midwives are important figures in promoting VBAC in suitable women.

There is a three year gap between the two groups of [women](#) but in terms of age, socio-economics, ethnic group and other variables, the groups are similar. The only significant change was the change in policy to midwife led care in the hospital. The results are also unlikely to be due to cultural changes over time as the VBAC rates across England more generally did not change during the study period.

Dr White added: "The more we understand the role of the microbes living in the vagina, the more we are uncovering the protective effects of labour and natural birth against conditions such as diabetes and obesity.

"Midwife led care is cost-effective and, if it safely reduces the number of repeat caesarean sections, then it's even more beneficial for mothers and babies."

More information: Helen K. White et al.
Evaluating a Midwife-Led Model of Antenatal Care for Women with a Previous Cesarean Section: A Retrospective, Comparative Cohort Study, *Birth* (2016). [DOI: 10.1111/birt.12229](https://doi.org/10.1111/birt.12229)

Provided by University of Manchester

APA citation: Natural births after caesarean more likely if you call the midwife (2016, April 20) retrieved 3 May 2021 from <https://medicalxpress.com/news/2016-04-natural-births-caesarean-midwife.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.