

# The effects of exercise on colorectal cancer survivors

26 April 2016



James Devin has studied the effects of exercise on colorectal cancer survivors.

A researcher who was born in the year that actress Audrey Hepburn and footballer Bobby Moore died of colorectal cancer is helping to lead recovery from the deadly condition.

University of Queensland researcher James Devin, 23, of the School of Human Movement and Nutrition Sciences, has studied the effects of exercise on colorectal cancer survivors.

"Increases in cardiorespiratory fitness are associated with a significant reduction in cancer-specific mortality," Mr Devin said.

"We showed that eight weeks of high-intensity interval training (HIIT) and moderate-intensity exercise (MIE) promoted significant improvements in [cardiorespiratory fitness](#) in colorectal cancer survivors.

"While the comparative effectiveness was not too different, HIIT may offer more sustained improvements following a detraining period.

"Encouragingly, no severe adverse events

occurred as a result of either types of training."

Mr Devin's research scored him the runner-up spot for the Young Investigator Award at the 2016 Exercise and Sports Science Australia (ESSA) Research to Practice Conference in Melbourne.

He collaborated with UQ's Associate Professor David Jenkins, Dr Kate Bolam, Dr Tina Skinner, Andrew Sax and Gareth Hughes.

The team monitored 35 colorectal [cancer](#) survivors as they undertook three sessions of exercise a week using either HIIT or MIE methods.

HIIT consisted of four rounds of four minutes of [exercise](#) at 85-95 per cent of maximum heart rate, and MIE consisted of 50 minutes of work at 70 per cent of the [maximum heart rate](#).

"Those who undertook HIIT significantly improved and maintained a 4.2 millilitres per kilogram per minute increase in cardiorespiratory output even after four weeks of detraining," Mr Devin said.

"The MIE group also significantly improved, but lost gains more rapidly over the detraining period.

"To put it in perspective, a 3.5 millilitres per kilogram per minute increase is associated with a 10 per cent reduction in mortality risk, so the improvements we have observed appear to be quite clinically significant for survivors."

Results of the research are published in the *Journal of Cancer Survivorship*.

**More information:** James L. Devin et al. The influence of high-intensity compared with moderate-intensity exercise training on cardiorespiratory fitness and body composition in colorectal cancer survivors: a randomised controlled trial, *Journal of Cancer Survivorship* (2015). [DOI: 10.1007/s11764-015-0490-7](#)

Provided by University of Queensland

APA citation: The effects of exercise on colorectal cancer survivors (2016, April 26) retrieved 30 August 2022 from <https://medicalxpress.com/news/2016-04-effects-colorectal-cancer-survivors.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*