

Prevalence of gestational diabetes varies across seasons

May 12 2016



(HealthDay)—The prevalence of gestational diabetes mellitus (GDM)

varies across the seasons, with higher prevalence in the summer and lower prevalence in the winter, according to a study published online May 10 in *Diabetes Care*.

Robert G. Moses, M.D., from Wollongong Hospital in Australia, and colleagues examined the effect of different seasons on the prevalence of GDM. All pregnancy glucose tolerance tests (GTTs) were collected over a three-year period in a temperate climate; results were available for 7,369 pregnancy GTTs.

The researchers found that the median one- and two-hour glucose results after GTT were significantly lower in winter than the overall one- and two-hour results (P

"The prevalence of GDM varies according to seasons, which leads to the possible overdiagnosis of GDM in [summer](#) and/or underdiagnosis in [winter](#)," the authors write. "Further research into standardization of the GTT or seasonal adjustment of the results may need to be considered."

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

Citation: Prevalence of gestational diabetes varies across seasons (2016, May 12) retrieved 5 February 2024 from <https://medicalxpress.com/news/2016-05-prevalence-gestational-diabetes-varies-seasons.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--