

Researchers determine the best strategy for preventing ulcers when taking NSAIDs

May 13 2016

Non-steroidal anti-inflammatory drugs (NSAIDs)—including ibuprofen, diclofenac, naproxen and others—are commonly used pain medications that are generally safe but may increase the risk of developing stomach and intestinal ulcers.

After researchers analyzed a large number of [clinical trials](#) that compared different ways of reducing these risks of NSAIDs, they found that the best strategy with the lowest overall risk was to combine a certain type of NSAID, known as a COX-2-selective NSAID, with a [proton pump inhibitor](#) (PPI). PPIs are most often used to treat heartburn and gastro-oesophageal reflux disease.

"The combination of a COX-2-selective NSAID with a PPI will be expensive and is not recommended for all patients who need to be on a NSAID; however, it is the safest and most effective treatment strategy for those at high risk of ulcer bleeding from NSAID treatment," said Prof. Jin Ling Tang, co-author of the Alimentary Pharmacology and Therapeutics study.

More information: J.Q.Yuan, et al. Systematic review with network meta-analysis: comparative effectiveness and safety of strategies for preventing NSAID-associated gastrointestinal toxicity. [DOI: 10.1111/apt.13642](#)

Provided by Wiley

Citation: Researchers determine the best strategy for preventing ulcers when taking NSAIDs (2016, May 13) retrieved 5 March 2023 from <https://medicalxpress.com/news/2016-05-strategy-ulcers-nsaids.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.