

Pressure recording analytical method estimates cardiac output

6 September 2016



agreement with the Fick method measurement," the authors write.

One author disclosed financial ties to Vygon.

More information: <u>Abstract</u> <u>Full Text (subscription or payment may be required)</u>

Copyright © 2016 HealthDay. All rights reserved.

(HealthDay)—The MostCare pressure recording analytical method (PRAM) is comparable to the Fick method of cardiac output estimation in pediatric patients, according to a study published online Aug. 27 in *Pediatric Anesthesia*.

José M. Alonso-Iñigo, M.D., Ph.D., from Hospital Clínico Universitario in Valencia, Spain, and colleagues conducted a single-center observational study involving 52 <u>pediatric patients</u> who underwent diagnostic right and left catheterization. Under hemodynamically stable conditions, cardiac index (CI) measurements were obtained with the MostCare-PRAM and Fick methods.

The researchers observed good agreement for $\rm CI_{Fick}$ and $\rm CI_{PRAM}$ in the 40 CI measurements (r² = 0.90; P

"In pediatric patients undergoing diagnostic right and left heart catheterization, the MostCare-PRAM was shown to estimate CI with a good level of



APA citation: Pressure recording analytical method estimates cardiac output (2016, September 6) retrieved 11 October 2022 from <u>https://medicalxpress.com/news/2016-09-pressure-analytical-method-cardiac-output.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.