

Study looks at the influence of fat when gut bacteria is reduced by antibiotics

8 September 2016, by Angela Koenig



Mary Emery Professor of Pathology and Laboratory Medicine at the UC College of Medicine. Credit: University of Cincinnati

A study led by University of Cincinnati (UC) lipid metabolism researchers lends additional insight into how bacteria in the gut, or lack thereof, influences intestinal mast cells (MMC) activation and perhaps fat absorption. The study, "Antibiotics Suppress Activation of Intestinal Mucosal Mast Cells and Reduce Dietary Lipid Absorption in Sprague-Dawley Rats," is currently available in the online edition of *Gastroenterology*.

"A change in the absorption of fat is very important for the development of obesity studies," says the study's principal investigator Patrick Tso, PhD, Mary Emery Professor of Pathology and Laboratory Medicine at the UC College of Medicine. According to the National Institute of Diabetes and Digestive and Kidney Diseases of the National Institutes of Health (NIH), overweight and obesity are risk factors for Type 2 diabetes, heart disease, high blood pressure, and other health problems. In the study, funded by NIH R01 DK092138, Tso and his team compared intestinal fat absorption and mast cell activation in rodents placed on antibiotics to clear their systems of bacteria and a control group, and then introduced lipids (fats) to both groups.

"Our results suggest that the presence of gut microbiota is involved in MMC activation, induced by the ingestion of fat, and contributes to fat-induced permeability," says Tso, adding that the

study also found a "novel role of the gut microbiome in the promoting the absorption of lipid."

Mast cells are intimately involved with the immune system in the body. They come from bone marrow and go into all tissues of the body. Each mast cell contains secretory granules (storage sacs), each containing powerful biologically active molecules called mediators. These can be secreted when [mast cells](#) are activated, leading to allergic and inflammatory responses.

Mast cells also play a key role in the defense of the gastrointestinal tract against foreign organisms, says Tso, and while the absorption of fat is actually associated with mucosal mast cell activation, it is not known exactly what role gut bacteria plays.

"This effect is specific to [fat](#), since the [absorption](#) of carbohydrates or protein does not activate mucosal mast cells," says Tso.

Provided by University of Cincinnati Academic Health Center

APA citation: Study looks at the influence of fat when gut bacteria is reduced by antibiotics (2016, September 8) retrieved 20 November 2022 from <https://medicalxpress.com/news/2016-09-fat-gut-bacteria-antibiotics.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.