

Bilateral tinnitus is hereditary

9 March 2017

Researchers have been able to demonstrate the hereditary nature of certain forms of tinnitus. Bilateral tinnitus - that is, tinnitus in both ears - has been shown to depend on genetic factors, particularly in men. The twin study, which is published in the journal *Genetics in Medicine*, was conducted by researchers at Karolinska Institutet together with colleagues from the European research network TINNET.

Ringling in the ears, a condition called [tinnitus](#), is experienced by 15 per cent of people in Sweden as well as in Europe. For one or two per cent of the population, the symptoms are extremely distressing and impact adversely on daily activities, work and sleep. Tinnitus thus has negative social consequences for the sufferers, while being an important economic burden to society.

Tinnitus prevalence increases with age and is thought to be related to a number of environmental factors but little research has been done on the subject. There are also no effective cures for the condition, due possibly to the heterogeneity of the condition.

Using data from the Swedish Twin Registry, researchers at Karolinska Institutet have found evidence that in some cases tinnitus has genetic causes.

"We've been able to show that different forms of tinnitus have a significant heritability and thus a dominant genetic influence over environmental factors," says Christopher R. Cederroth at Karolinska Institutet's Department of Physiology and Pharmacology.

When the researchers first examined all forms of tinnitus they made the same conclusions on heritability as others have reported. It was only after grouping the subjects by sex and unilateral/bilateral tinnitus that they uncovered the genetic correlation.

"This result is surprising and unexpected as it

shows that, unlike the conventional view of tinnitus being driven by environmental factors, there is a genetic influence for bilateral tinnitus which is more pronounced in men" says Dr Cederroth.

Their discovery also shows that bilateral and unilateral tinnitus constitute two separate sub-groups, only one of which is influenced by [genetic factors](#). This, claims Dr Cederroth, not only has considerable clinical relevance but is also important from a public health perspective:

"Tinnitus sufferers need better care and treatment than they're currently getting. We need more genetic studies and a better molecular understanding of its generation, which could open unforeseen avenues to drug development."

More information: Genetic susceptibility to bilateral tinnitus in a Swedish twin cohort", Iris Lianne Maas, Petra Bru?ggemann, Teresa Requena Navarro, Jan Bulla, Niklas K. Edvall, Jacob v.B. Hjelmberg, Agnieszka J. Szczepek, Barbara Canlon, Birgit Mazurek, Jose A. Lopez-Escamez, Christopher R. Cederroth. *Genetics in Medicine*, online 9 March 2017, [DOI: 10.1038/GIM.2017.4](#)

Provided by Karolinska Institutet

APA citation: Bilateral tinnitus is hereditary (2017, March 9) retrieved 18 August 2022 from <https://medicalxpress.com/news/2017-03-bilateral-tinnitus-hereditary.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.