


Conscious sedation is a safe alternative to general anesthesia for heart valve procedure

11 April 2017



Rendering of a heart valve inserted during transcatheter aortic valve replacement. A UCLA study found that keeping patients awake but sleepy during the procedure is preferable to general anesthesia. Credit: Edwards Lifesciences

UCLA scientists have found that conscious sedation—a type of anesthesia in which patients remain awake but are sleepy and pain-free—is a safe and viable option to general anesthesia for people undergoing a minimally invasive heart procedure called transcatheter aortic valve replacement.

In the study, patients who underwent conscious sedation had a similar rate of adverse events to those who underwent anesthesia, but those who were given conscious sedation had shorter stays in the [intensive care unit](#) (30 versus 96 hours for those with general anesthesia) and shorter hospital stays (4.9 days versus 10.4 days). The direct costs for their care were 28 percent lower, and almost all other medical costs were lower as well. In both groups, 1.5 percent of patients died during hospitalization.

Aortic stenosis is a common disease, particularly among people 75 and older, in which one of the valves of the heart narrows and impedes blood flow to the body. Previously, treating [aortic stenosis](#) required [open heart surgery](#), which is particularly risky for elderly and frail patients.

In recent years, the use of transcatheter aortic valve replacement, or TAVR, has made the repair safer: Physicians place a new valve in the heart via a catheter inserted through the patient's leg, which results in less injury and an easier recovery than open heart surgery. Most medical centers perform the procedure under general anesthesia but some, including UCLA, now offer their patients conscious sedation, which does not require a breathing tube.

The study was the first to detail the cost differences and outcomes between conscious sedation and general anesthesia during transcatheter aortic valve replacement. Researchers analyzed records of 196 adult patients who underwent transcatheter aortic valve replacement at Ronald Reagan UCLA Medical Center between August 2012 and June 2016. The researchers write that a large-scale, controlled and randomized study should be conducted to validate their findings.

Based on the study, patients undergoing transcatheter [aortic valve replacement](#) should receive [conscious sedation](#) instead of [general anesthesia](#) whenever possible.

More information: William Toppen et al. Improved costs and outcomes with conscious sedation vs general anesthesia in TAVR patients: Time to wake up?, *PLOS ONE* (2017). [DOI: 10.1371/journal.pone.0173777](#)

Provided by University of California, Los Angeles

APA citation: Conscious sedation is a safe alternative to general anesthesia for heart valve procedure (2017, April 11) retrieved 27 April 2021 from <https://medicalxpress.com/news/2017-04-conscious-sedation-safe-alternative-anesthesia.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.