

Tidal CO₂ prognostic for chronic thromboembolic pulmonary HTN

28 August 2017



class were seen for patients with inoperable CTEPH who had ETCO₂

(HealthDay)—End-tidal carbon dioxide (ETCO₂) levels are prognostic for patients with chronic thromboembolic pulmonary hypertension (CTEPH), according to a letter to the editor published online Aug. 14 in the *Annals of the American Thoracic Society*.

Manuel J. Richter, M.D., from the Kerckhoff Heart and Thoracic Center in Bad Nauheim, Germany, and colleagues retrospectively analyzed data from the Giessen Pulmonary Hypertension Registry for 151 patients with inoperable CTEPH. Baseline ETCO₂ levels obtained at rest were available for all patients.

The researchers found that 27.2 percent of the patients died during a mean follow-up of 64.8 ± 40.2 months. The median ETCO₂ at baseline was 27 mm Hg, which was substantially decreased. ETCO₂ was related to overall survival as a continuous variable in univariate analysis (hazard ratio, 0.91). Significantly impaired pulmonary hemodynamics and a more advanced functional

APA citation: Tidal CO2 prognostic for chronic thromboembolic pulmonary HTN (2017, August 28)
retrieved 1 July 2022 from <https://medicalxpress.com/news/2017-08-tidal-cosub2sub-prognostic-chronic-thromboembolic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.