

Physical activity can lead to difference in diet preferences between males, females

5 September 2017, by Jeff Sossamon



Credit: CC0 Public Domain

Recent studies have shown that approximately 90 percent of adult Americans fail to reach the U.S. Department of Health guidelines for physical activity, which could be contributing to surging obesity rates. Now, new research by a multidisciplinary team of University of Missouri researchers suggests that physical activity can change diet preferences in males, but not in females—an area that researchers say has not been thoroughly studied.

"Our team wanted to make every effort to study female perspectives on how exercise affects [diet](#), because most other studies neglect females," said Jenna Lee, a doctoral candidate in the Interdisciplinary Neuroscience Program at MU. "We wanted to take a look at what drives diet preference and if environmental factors, such as [physical activity](#), play a role in how males and females eat."

The team, led by Matthew Will, an associate professor of psychological sciences in the MU College of Arts and Science, divided male and [female rats](#) into two groups—a sedentary group and

one that had access to a running wheel. Both groups ate the same food. After another week, Lee replaced the standard diet with three optional diets: high fat (similar to cookie dough), high sucrose (three times more sucrose than the other two diets), and a high-cornstarch diet. Each of the diets were matched on proteins, and the rats had continuous access to all three diets for four weeks.

Results showed that sedentary male rats preferred the high-fat diet over the other diets. Male runners ate about half as much of the high-fat diet as their sedentary counterparts but increased their intake of the other two options. Female sedentary rats, like their male counterparts, mostly stuck to the high-fat diet. Surprisingly, female runners also preferred the high-fat diet and actually consumed slightly more calories than the sedentary females.

"We also examined brain opioids and [gut microbiota](#), and we discovered key changes that paralleled the patterns observed in diet preferences between male and female runners. A reason for this might be that females have an elevated threshold for rewards. Considering females demonstrate higher levels of reward signaling in the brain, this may possibly explain the higher threshold or capacity for reward. Perhaps something like running may be satiating for males but not for the females, so the females are consuming more of the [high-fat diet](#). We expected to find differences between runners and sedentary [rats](#), but it was the sex differences that surprised us."

The research draws attention to the importance of studying both men and women in research, Lee said.

The study, "Sex determines effect of physical activity on diet preference: Association of striatal opioids and gut microbiota composition," was published by *Behavioral Brain Research*.

More information: Jenna R. Lee et al. Sex determines effect of physical activity on diet preference: Association of striatal opioids and gut microbiota composition, *Behavioural Brain Research* (2017). [DOI: 10.1016/j.bbr.2017.07.018](https://doi.org/10.1016/j.bbr.2017.07.018)

Provided by University of Missouri-Columbia

APA citation: Physical activity can lead to difference in diet preferences between males, females (2017, September 5) retrieved 17 August 2022 from <https://medicalxpress.com/news/2017-09-physical-difference-diet-males-females.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.