


Discontinuity of care puts older patients at risk of emergency hospital admission

14 November 2017



Credit: University of Bristol

Older patients who do not see the same GP over a period of time are at higher risk of emergency hospital admission than those who see the same or a small number of GPs, according to new research.

Researchers at the Universities of Bristol and Oxford found that the risk of [emergency hospital admission](#) was more than twice as high for patients with the least continuity of care compared with those with high continuity of care.

The study, published in *Annals of Family Medicine* today, involved analysing data from 10,000 electronic records of patients aged 65 years and older, from 297 general practices in England, between April 2010 and March 2014. These data were linked to hospital records to measure the association between continuity of care and risk of emergency hospital admission. The team focused on [older patients](#) as they are seen more frequently in general practice than younger adult patients and are most at risk of emergency hospital admission.

Acute hospital services in England are under

sustained pressure with increasing emergency attendances, resulting in longer waiting times in emergency departments and high bed occupancy rates. These are both associated with poorer clinical outcomes as well as greater demands on staff. Previous evidence suggests that some patients admitted as emergencies could be managed at home. Initiatives to enhance continuity of care, the researchers suggest, could potentially reduce hospital admissions.

Dr Peter Tammes, Senior Research Associate at the University of Bristol's Centre for Academic Primary Care and lead author said: "Our study shows that lack of continuity of care in this high-risk age group is associated with more frequent emergency admissions, which in turn suggests that better continuity of care might lead to a reduction in admissions. Discontinuity of care reduces the opportunity for building trust and mutual responsibility between doctors and patients, which might underlie the increased risk of emergency hospital [admission](#)."

"More research is needed to have a clearer understanding of how this association works and to understand patients' values and experiences. It would also be helpful to evaluate new schemes to improve continuity of care, such as the introduction in 2014 of a named GP for elderly patients - especially as the merging of practices into 'super-practices' is expected to lead to an overall decrease in continuity of care.

"Despite the challenges, we hope that these findings support policy makers, commissioners, [patients](#) and GPs in their efforts to improve continuity of care, where the evidence supports it."

More information: Peter Tammes et al. Continuity of Primary Care and Emergency Hospital Admissions Among Older Patients in England, *The Annals of Family Medicine* (2017). [DOI: 10.1370/afm.2136](#)

Provided by University of Bristol

APA citation: Discontinuity of care puts older patients at risk of emergency hospital admission (2017, November 14) retrieved 15 November 2022 from <https://medicalxpress.com/news/2017-11-discontinuity-older-patients-emergency-hospital.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.