

Postnatal depression has greater impact on children's development when it is persistent and severe

1 February 2018



Postnatal depression which persists beyond 6 months after birth and is severe, increases the risk of children exhibiting behavioural problems, achieving lower GCSE mathematics grades at 16 years and having depression at 18 years of age. Postnatal depression which is persistent (whether moderate or severe) increases mothers' risk of continuing to experience depressive symptoms beyond the postnatal year, with high levels found up until 11 years after childbirth.

Available research suggests that [postnatal depression](#) is associated with increased risks to [children's development](#) affecting a range of domains. The effects are variable with some evidence suggesting that brief episodes, while distressing to the mother, may not impact negatively on children's development. However, episodes of [depression](#) which persist for six months or more in duration may increase the risk for children. Identifying women at most risk is important both for women's mental health and

children's development. The authors investigated whether depression has a similar impact when it is, or is not, persistent at either moderate or severe levels. Researchers from the Universities of Oxford, Bristol, Reading and UCLA in a new paper in JAMA Psychiatry tracked changes in [mothers' depressive symptoms](#) following the postnatal year and differences in children's development when postnatal depression was, or was not, persistent.

The sample consisted of 9848 mothers and 8419 children who were taking part in the Avon Longitudinal Study of Parents and Children (ALSPAC). The study utilises data from different stages of children's development ranging from preschool to late adolescence; including behaviour problems at 3.5 years of age, GCSE mathematics grades at 16 years of age and depression at 18 years of age to determine potential effects both in the short and long term.

The study highlights for the first time that when postnatal depression is both persistent and severe the risk of negative outcomes to children's development is raised. Furthermore, when depression persists beyond the first 6 months after birth, the risk for mothers continuing to experience depressive symptoms beyond the postnatal year is also increased. The authors report that these symptoms can continue up to 11 years after childbirth, but it is possible that they persist even longer. For two of the outcomes (GCSE mathematics grades at 16 and depression at 18 years), when postnatal depression either did not persist, or persisted only at moderate severity, the odds of achieving lower GCSE mathematics grades or experiencing depression at 18 years were similar to those of children whose mothers did not experience depressive symptoms at all in the postnatal year. This suggests that long term negative effects on mathematics grades and

depression in adolescence principally occur in the context of [persistent depression](#), but not with briefer and/or less severe episodes.

The study identifies a group of mothers and children who should be prioritised for treatment both to address [maternal depression](#) that is more likely to continue, and to mitigate the effects on children's development. The authors have recently shown that treatment for women with persistent postnatal depression can be delivered effectively with high rates of sustained remission using home-based delivery of a psychological therapy.

Dr. Elena Netsi from the Department of Psychiatry at Oxford University and lead author on the paper said: "Depression which persists beyond the first 6 months after birth should alert health care professionals to a depression which may become chronic. While screening for postnatal mental health problems in the UK already occurs within the first 3 months, screening once more in the second part of the postnatal year will help identify families who should be prioritised for treatment, both to address the mother's depression and to mitigate potential effects on children's development. Given that there are effective psychological therapies for depression it is important that help is provided for these women."

More information: Elena Netsi et al. Association of Persistent and Severe Postnatal Depression With Child Outcomes, *JAMA Psychiatry* (2018).
[DOI: 10.1001/jamapsychiatry.2017.4363](https://doi.org/10.1001/jamapsychiatry.2017.4363)

Provided by University of Bristol

APA citation: Postnatal depression has greater impact on children's development when it is persistent and severe (2018, February 1) retrieved 30 April 2021 from <https://medicalxpress.com/news/2018-02-postnatal-depression-greater-impact-children.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.