

Many epilepsy patients take drug combinations that interact

7 February 2018



availability of newer drugs," the authors wrote.

More information: Edward Faught et al, Risk of pharmacokinetic interactions between antiepileptic and other drugs in older persons and factors associated with risk, *Epilepsia* (2018). [DOI: 10.1111/epi.14010](https://doi.org/10.1111/epi.14010)

Provided by Wiley

Generalized 3 Hz spike and wave discharges in a child with childhood absence epilepsy. Credit: Wikipedia.

In an *Epilepsia* analysis of 2008-2010 Medicare claims data, one in four older Americans with new-onset epilepsy and more than one-third with prevalent epilepsy received a combination of antiepileptic drugs and non-epilepsy drugs that could interact to alter the effectiveness of the non-epilepsy drugs. Also, more than 1 in 5 patients received a drug combination that could alter the effect of the antiepileptic drugs and potentially cause toxicity.

Having interacting drug pair combinations was more likely for those with [comorbid conditions](#) and those eligible for a low-income subsidy.

"Avoidance of interacting [antiepileptic drugs](#) is feasible for most persons because of the

APA citation: Many epilepsy patients take drug combinations that interact (2018, February 7) retrieved 12 October 2022 from

<https://medicalxpress.com/news/2018-02-epilepsy-patients-drug-combinations-interact.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.