

# Choroidal thickness changes in patients with untreated DM

April 23 2018

---



(HealthDay)—The choroidal thickness changes in patients with

untreated diabetes mellitus over time, according to a study published online April 14 in *Clinical & Experimental Ophthalmology*.

Hiroaki Endo, from Teine Keijinkai Hospital in Sapporo, Japan, and colleagues examined the choroidal [layer](#) thickness in 268 eyes of 134 patients with diabetes mellitus and 72 healthy controls (92 eyes) to compare the findings based on medical history of systemic diabetes mellitus treatments and stage of diabetic retinopathy (DR).

The researchers observed a strong correlation between masked raters of choroidal thickness measurements, proving high reproducibility. In the untreated diabetes mellitus group, the total and outer choroid thicknesses in mild/moderate non-proliferative DR (mNPDR) were significantly thinner versus normal controls. In severe NPDR, choroidal outer layer thickness in the untreated group was significantly thicker than normal controls. However, in the group treated for [diabetes](#) mellitus, there were no significant differences regarding choroidal layer thicknesses and all stages of DR.

"The choroidal thickness significantly changed in the [diabetes mellitus](#) untreated group, and the main anatomical changes might result from the outer layer," the authors write.

**More information:** [Abstract/Full Text \(subscription or payment may be required\)](#)

Copyright © 2018 [HealthDay](#). All rights reserved.

Citation: Choroidal thickness changes in patients with untreated DM (2018, April 23) retrieved 25 December 2022 from <https://medicalxpress.com/news/2018-04-choroidal-thickness-patients-untreated-dm.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.