

Weight loss surgery may cause significant skeletal health problems

2 May 2018

A new JBMR Plus review examines the negative impacts of weight loss surgery on bone health.

The review of published studies notes that weight loss surgery can cause declines in bone mass and strength, and it is linked with an increased risk of bone fractures. Skeletal changes after surgery appear early and continue even after weight loss plateaus and weight stabilizes. Nutritional factors, mechanical unloading, hormonal factors, and changes in body composition and bone marrow fat may contribute to poor bone health.

Most studies have examined the effects of the Roux-en-Y gastric bypass procedure, which was the most commonly performed weight loss procedure worldwide until it was very recently overtaken by sleeve gastrectomy. Because sleeve gastrectomy is a newer procedure, its skeletal effects have not yet been well defined.

The review's findings indicate that clinical guidelines on weight loss surgery should address bone health as a priority. "Current clinical guidelines do address bone health, but most recommendations are based on low-quality evidence or expert opinion," said co-author Dr. Anne Schafer, of the University of California, San Francisco and the San Francisco VA Health Care System. "Future studies should address strategies to avoid long-term skeletal consequences of these otherwise beneficial procedures."

More information: Claudia Gagnon et al, Bone Health After Bariatric Surgery, JBMR Plus (2018). DOI: 10.1002/jbm4.10048

Provided by Wiley APA citation: Weight loss surgery may cause significant skeletal health problems (2018, May 2) retrieved 11 October 2022 from

https://medicalxpress.com/news/2018-05-weight-loss-surgery-significant-skeletal.html



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.