

# Artery hardening and thickness not affected by stopping hormone therapy

18 July 2018

Heart disease is still the number one killer of US women, and hormone therapy remains a top treatment for menopause symptoms. A new study connects these two facts to demonstrate little effect of hormone therapy on artery thickness as a precursor to heart disease. Study results are published online today in *Menopause*, the journal of The North American Menopause Society (NAMS).

According to the American Heart Association, [heart disease](#) is responsible for one in three deaths of US women each year. That means about one woman dies every minute. Statistics like these drive a continued interest in the medical field to study risk factors that may affect the incidence of heart disease. This includes the ongoing debate regarding the effect of [hormone therapy](#) on the progression of cardiovascular disease.

The well-known Kronos Early Estrogen Prevention Study (KEEPS) was designed to compare the effects of lower doses of either oral conjugated equine estrogens (CEE) or transdermal estradiol on the progression of atherosclerosis (hardening and thickening of the arteries), the most common cause of heart attacks and strokes. After 4 years of therapy, no differences in artery thickness were detected among the oral CEE-, transdermal estradiol-, and placebo-treated groups. In this new study, a subgroup of KEEPS participants were further evaluated for 3 years after the initial study.

The results of the follow-up study appear in the article "Changes in carotid artery intima-media thickness 3 years after cessation of [menopausal hormone therapy](#): follow-up from the Kronos Early Estrogen Prevention Study." Study results demonstrate that the cessation of hormone therapy at the lower doses did not accelerate changes in the thickness of the arteries.

"In the KEEPS trial, low doses of hormone therapy (oral or transdermal) given for 4 years close to

menopause did not improve carotid artery intima-media thickness (CIMT), and this study shows no rebound effect on progression of CIMT after stopping the hormones," says Dr. JoAnn Pinkerton, executive director of NAMS. "In contrast, the Early Versus Late Intervention Trial With Estradiol (ELITE) found benefits on CIMT when hormone therapy was given early but not when started more than 10 years from menopause. The primary indication for the use of [hormone therapy](#) should be for relief of bothersome [menopause symptoms](#) or for those at elevated risk of bone loss. There may be beneficial effects on reduction of [heart disease](#), but it should not be used for its prevention."

Provided by The North American Menopause Society

APA citation: Artery hardening and thickness not affected by stopping hormone therapy (2018, July 18) retrieved 15 September 2022 from <https://medicalxpress.com/news/2018-07-artery-hardening-thickness-affected-hormone.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*