

Smoking during pregnancy may lead to childhood eye condition

7 November 2018

Provided by Wiley



Credit: CC0 Public Domain

In an *Acta Ophthalmologica* analysis of 11 relevant articles, maternal smoking during pregnancy was associated with a 46 percent increased risk that offspring will develop strabismus—one of the most prevalent eye-related diseases among children. Maternal smoking of ?10 cigarettes per day during pregnancy was linked with a 79 percent increased risk of strabismus in offspring.

In children with strabismus, the eyes do not properly align with each other. This can contribute to [visual impairment](#) and emotional problems.

"Maternal smoking during pregnancy is an important public health problem, particularly in developed countries, and its effect on offspring eye health deserves our attention," said senior author Dr. Zuxun Lu, of the Huazhong University of Science and Technology, in China.

More information: Yudi Yang et al, Maternal smoking during pregnancy and the risk of strabismus in offspring: a meta-analysis, *Acta Ophthalmologica* (2018). DOI: [10.1111/aos.13953](https://doi.org/10.1111/aos.13953)

APA citation: Smoking during pregnancy may lead to childhood eye condition (2018, November 7) retrieved 2 September 2022 from <https://medicalxpress.com/news/2018-11-pregnancy-childhood-eye-condition.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.