

Increased risk of comorbid neck injury in females with a concussion-related ED visit

January 7 2019



Credit: Mary Ann Liebert, Inc., publishers

In a study of neck injury comorbidity in concussion-related emergency department (ED) visits in Ontario, Canada, females had significantly higher odds than males of sustaining a comorbid neck injury in a multitude of circumstances and stages of their lives. Sex differences in the rate of neck injury comorbidity among patients with a concussion-related ED visit are reported in an article published in *Journal of Women's Health*.

In the article, "Neck Injury Comorbidity in Concussion-Related Emergency Department Visits: A Population-Based Study of Sex Differences Across the Lifespan," the researchers highlight the important influences of sex and age on neck injury comorbidity in concussion-related ED visits. Data on ED visits in Ontario were extracted from the Canadian Institute for Health Information National Ambulatory Care Reporting System. The International Classification of Diseases, 10th Revision, code for concussion was used to identify patients with a first concussion-related ED visit between fiscal years 2002/03 and 2011/12 (inclusive). The article is coauthored by Mitchell Sutton, MSc and colleagues from the Toronto Rehabilitation Institute-University Health Network, the University of Toronto, and the Institute for Clinical Evaluative Sciences (Toronto, Canada).

"The finding that females with a concussion-related ED visit have an increased risk of sustaining a comorbid neck [injury](#) has important clinical implications," states Susan G. Kornstein, MD, Editor-in-Chief of *Journal of Women's Health* and Executive Director of the Virginia Commonwealth University Institute for Women's Health, Richmond, VA. "In the ED setting, increased consideration of potential comorbid [neck](#) injuries, especially among females ages 5 to 49 years, is warranted to enable timely intervention and treatment."

More information: Mitchell Sutton et al, Neck Injury Comorbidity in Concussion-Related Emergency Department Visits: A Population-Based

Study of Sex Differences Across the Life Span, *Journal of Women's Health* (2018). [DOI: 10.1089/jwh.2018.7282](https://doi.org/10.1089/jwh.2018.7282)

Provided by Mary Ann Liebert, Inc

Citation: Increased risk of comorbid neck injury in females with a concussion-related ED visit (2019, January 7) retrieved 24 November 2023 from <https://medicalxpress.com/news/2019-01-comorbid-neck-injury-females-concussion-related.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.