

Acupressure reduces lasting symptoms in breast cancer survivors

13 March 2019



acupressure was associated with greater reductions in pain interference. There were no statistically significant moderators of sleep quality, anxiety, or depressive symptoms.

"Acupressure was associated with greater improvements than usual care in anxiety, pain, and symptoms of depression in breast cancer survivors with troublesome fatigue," the authors write.

More information: Abstract/Full Text

Copyright © 2019 HealthDay. All rights reserved.

(HealthDay)—Acupressure is associated with improvements in co-occurring symptoms in breast cancer survivors with fatigue, according to a study recently published in *JNCI Cancer Spectrum*.

Suzanna Maria Zick, M.P.H., from the University of Michigan in Ann Arbor, and colleagues evaluated changes in <u>depressive symptoms</u>, anxiety, and <u>pain</u> before and after randomization to selfacupressure (either relaxing or stimulating) or usual care (typical sleep-management techniques) among 288 <u>breast cancer survivors</u> with fatigue participating in a clinical trial.

The researchers found that after treatment, depressive symptoms improved significantly for the relaxing acupressure group (41.5 percent) versus the stimulating acupressure (25 percent) and usual care (7.7 percent) groups. For anxiety, both acupressure groups were associated with greater improvements than usual care. Only relaxing acupressure was associated with greater reductions in <u>pain severity</u>, while only stimulating



APA citation: Acupressure reduces lasting symptoms in breast cancer survivors (2019, March 13) retrieved 13 June 2022 from <u>https://medicalxpress.com/news/2019-03-acupressure-symptoms-breast-cancer-survivors.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.