


Drug reduces risk of kidney failure in people with diabetes, study finds

15 April 2019, by Amy Jeter Hansen



This image shows a cross section of a kidney. Credit: Holly Fischer/Wikipedia

A new landmark clinical trial shows that a drug lowers the risk of kidney failure by a third in people with Type 2 diabetes and kidney disease.

"For the first time in 18 years, we have a therapy for [patients](#) with Type 2 [diabetes](#) and chronic [kidney disease](#) that decreases [kidney failure](#)," said Kenneth Mahaffey, MD, professor of medicine at the School of Medicine and co-principal investigator of the trial. "Now, patients with diabetes have a promising option to guard against one of the most severe risks of their condition."

The trial involved 4,401 participants in 34 countries.

The drug, canagliflozin, improves on a nearly two-decades-old therapy that is currently the only treatment approved to protect [kidney function](#) in people with Type 2 diabetes. In the trial, canagliflozin also was found to reduce the risk of major cardiovascular events.

Canagliflozin increases the excretion of glucose through the kidneys. It has already been approved by the Food and Drug Administration to lower [blood glucose](#) in patients with Type 2 diabetes and to reduce the risk of major adverse cardiovascular events in patients with Type 2 diabetes and existing [heart disease](#).

A paper describing the findings of the CRENDENCE trial was published April 14 in the *New England Journal of Medicine* and presented at the International Society of Nephrology's World Congress of Nephrology in Melbourne. Mahaffey, who is director of the Stanford Center for Clinical Research, is the study's senior author. The lead author is Vlado Perkovic, MBBS, Ph.D., executive director of The George Institute for Global Health Australia, and a professor of medicine at the University of New South Wales in Sydney.

'Definitive trial result'

"People with diabetes and [kidney](#) disease are at extremely high risk of kidney failure, heart attack, stroke and death," Perkovic said. "With this definitive trial result, we now have a very effective way to reduce this risk using a once-daily pill."

Participants in the trial received the best care available for kidney disease under current guidelines, a type of therapy called renin-angiotensin-aldosterone system, or RAAS, blockade. In addition, half were randomly selected to receive canagliflozin, while the other half were

given a placebo.

The primary results of the study found that participants who took canagliflozin were 30 percent less likely than the placebo group to develop kidney failure or die from either [renal failure](#) or cardiovascular disease. Their risk of kidney failure or death from kidney failure was reduced by 34 percent, and the risk of hospitalization for heart failure or death due to cardiac causes decreased by 31 percent.

'Eagerly sought' treatment

People with diabetes can develop kidney disease because prolonged high blood sugar harms blood vessels in the kidney. In addition, diabetes often causes high blood pressure, which can stretch and weaken blood vessels in the organ.

For the past two decades, physicians have largely relied on RAAS blockade to prevent the deterioration of kidney function in diabetic patients. Although RAAS blockade lowers blood pressure and delays progression of kidney disease, patients undergoing this treatment remain at a high risk for renal failure and cardiovascular disease, as well as death from these conditions.

Given that the number of people with Type 2 diabetes worldwide is estimated to rise by 20 percent to 510 million in 2030, "a drug like canagliflozin that improves both cardiovascular and renal outcomes has been eagerly sought by both patients with Type 2 diabetes and clinicians caring for them," Mahaffey said.

More information: Vlado Perkovic et al. Canagliflozin and Renal Outcomes in Type 2 Diabetes and Nephropathy, *New England Journal of Medicine* (2019). [DOI: 10.1056/NEJMoa1811744](https://doi.org/10.1056/NEJMoa1811744)

Provided by Stanford University Medical Center
APA citation: Drug reduces risk of kidney failure in people with diabetes, study finds (2019, April 15) retrieved 12 October 2022 from <https://medicalxpress.com/news/2019-04-drug-kidney-failure-people-diabetes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.