

Between 16% and 18% of preadolescents have ideas of suicide

July 11 2019



The Universitat Roviri Researcher, Núria Voltas. Credit: URV

The researchers studied a group of 720 boys and 794 girls who attended 13 schools in Reus. They were monitored during three developmental periods: 10 years old, 11 years old and 13 years old. At the beginning of

the study, the students answered a series of psychological tests that were used to detect which of them presented emotional symptoms related to depression, anxiety and obsessive compulsive disorder (OCD). From their responses, two groups were created: one group at risk of emotional problems and a control group.

The disorders were diagnosed with standardised international criteria and the boys and girls were monitored to see how suicidal ideation developed throughout the research period.

The figures were quite stable. During the first period, 16% of the students stated that they had thought about suicide, of whom 33% stated the same one year later. In both the second and the third period, ideas of suicide were expressed by 18% of the students surveyed. The risk of suicide was determined in a personal interview and was present in 12.2% of the children with an average age of 11 years old. Although there were no differences between the sexes, the severity of the suicidal behaviour was greater in boys.

The researchers also observed what factors predicted suicidal ideation and they found here that there were differences between the sexes. "In boys it is previous depressive symptoms which determine subsequent [suicidal ideation](#)," says Núria Voltas, one of the researchers involved in the study. In girls, on the other hand, it is a combination of anxiety symptoms, OCD and the family's socioeconomic situation.

The results of this research, published in the scientific journal *Archives of Suicide Studies* reveal the factors that can trigger ideas of [suicide](#) in this age group. "Our results will enable us to have greater control over this particular aspect and take prevention measures in preadolescents, who are going through a [period](#) of considerable vulnerability," she concludes.

More information: Núria Voltas et al, Suicidality in a Community Sample of Early Adolescents: A Three-Phase Follow-Up Study, *Archives of Suicide Research* (2019). [DOI: 10.1080/13811118.2019.1588816](https://doi.org/10.1080/13811118.2019.1588816)

Provided by Universitat Rovira i Virgili

Citation: Between 16% and 18% of preadolescents have ideas of suicide (2019, July 11)
retrieved 7 May 2023 from
<https://medicalxpress.com/news/2019-07-preadolescents-ideas-suicide.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.