

One in ten older adults currently binge drinks

31 July 2019



Credit: CC0 Public Domain

More than a tenth of adults age 65 and older currently binge drink, putting them at risk for a range of health problems, according to a study by researchers at NYU School of Medicine and the Center for Drug Use and HIV/HCV Research (CDUHR) at NYU College of Global Public Health.

The study, published in the *Journal of the American Geriatrics Society*, also finds certain factors—including using cannabis and being male—are associated with an increase in [binge drinking](#).

Binge drinking is a risky behavior, particularly for [older adults](#) due to aging-related physical changes (for instance, an increased risk of falling) and the likelihood of having chronic health issues. Despite the potential for harm, little research has focused on binge drinking among older adults.

"Binge drinking, even episodically or infrequently, may negatively affect other health conditions by exacerbating disease, interacting with prescribed

medications, and complicating disease management," said Benjamin Han, MD, MPH, the study's lead author and an assistant professor in the Department of Medicine's Division of Geriatric Medicine and Palliative Care, and the Department of Population Health at NYU Langone Health.

In this study, Han and his colleagues used the most recent national data to determine the current prevalence and factors that may increase the risk of binge drinking among adults. The researchers examined data from 10,927 U.S. adults age 65 and older who participated in the National Survey on Drug Use and Health between 2015 and 2017. They looked at the prevalence of current (past-month) binge alcohol use, defined by the National Institute on Alcohol Abuse and Alcoholism (NIAAA) as five drinks or more on the same occasion for men and four drinks or more for women. They also compared demographic and health factors of past-month binge drinkers with people who drank within the past month, but below the binge drinking threshold.

The authors estimate that more than one in 10 (10.6 percent) older adults have binge drank in the past month—[an increase compared to earlier studies](#). In the decade leading up to the data used in this study (2005-2014), binge drinking among adults 65 and older was between 7.7 and 9 percent.

Binge drinkers were more likely to be male, current tobacco and/or cannabis users, African American, and have less than a high school education. They were also more likely to visit the emergency room in the past year. Similar to previous studies, the study did not find associations between binge drinking and other mental health disorders.

"The association of binge drinking with cannabis use has important health implications. Using both may lead to higher impairment effects. This is particularly important as cannabis use is becoming

more prevalent among older adults, and older adults may not be aware of the possible dangers of using cannabis with alcohol," said CDUHR researcher Joseph Palamar, Ph.D., MPH, the study's senior author and an associate professor in the Department of Population Health at NYU Langone Health.

Provided by New York University

The researchers also examined chronic disease profiles of older binge drinkers, and noted that binge drinkers had a lower prevalence of two or more chronic diseases compared to non-binge drinkers. The most common chronic disease among binge drinkers was hypertension (41.4 percent), followed by cardiovascular disease (23.1 percent) and diabetes (17.7 percent).

"Binge drinkers were less likely to have most chronic diseases compared to alcohol users who did not binge drink. This may be because some people stop or decrease their drinking when they have an illness or alcohol-related disease," said Han, who is also a CDUHR researcher. "Clinicians must be aware that some older adults with chronic disease still engage in binge drinking behaviors, which can worsen their [health](#) issues. This may explain why binge drinkers were more likely to report visits to the emergency room."

The researchers note that while the study uses the NIAAA's recommended threshold for binge drinking, the same organization also suggests lower drinking limits for adults over 65: no more than three drinks a day. Since the current analysis used the higher cutoff for binge drinking, the study may underestimate the prevalence of binge [drinking](#) among older adults.

"Our results underscore the importance of educating, screening, and intervening to prevent alcohol-related harms in older [adults](#), who may not be aware of their heightened risk for injuries and how alcohol can exacerbate [chronic diseases](#)," said Han.

More information: Benjamin H. Han et al, Binge Drinking Among Older Adults in the United States, 2015 to 2017, *Journal of the American Geriatrics Society* (2019). [DOI: 10.1111/jgs.16071](https://doi.org/10.1111/jgs.16071)

APA citation: One in ten older adults currently binge drinks (2019, July 31) retrieved 3 May 2021 from <https://medicalxpress.com/news/2019-07-ten-older-adults-binge.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.