


High-intensity step training boosts stroke survivors' walking skills

22 August 2019



Credit: CC0 Public Domain

High-intensity step training that mimics real-world conditions may better improve walking ability in stroke survivors compared to traditional, low-impact training, according to new research published in the American Heart Association's journal *Stroke*.

"People who suffer strokes often have difficulty walking and impaired balance. Rehabilitation after a [stroke](#) traditionally focuses on patients practicing low-intensity walking, usually only in a forward direction, which does not provide enough of a challenge to the [nervous system](#) to enable patients to negotiate real-world situations, such as uneven surfaces, stairs or changing direction," said study author T. George Hornby, Ph.D., professor of physical medicine and rehabilitation at Indiana University School of Medicine in Indianapolis. "Our study suggests that stroke patients can perform higher intensity walking exercises and more difficult tasks than previously thought possible. We need to move beyond traditional, low-intensity rehabilitation to challenge the nervous and cardiovascular systems so patients can improve function and perform better in the real world."

Researchers evaluated 90 people, 18- to 85-years-old with weakness on one side of the body who had survived a stroke at least six months prior.

Participants received training of either high-intensity stepping performing variable, [difficult tasks](#); high-intensity stepping performing only forward walking; or low-intensity stepping of variable tasks. Variable tasks included walking on uneven surfaces, up inclines and stairs, over randomly placed obstacles on a treadmill and across a balance beam.

The researchers found:

- Survivors in both the high-intensity, variable training and high-intensity, forward walking groups walked faster and farther than the low-intensity, variable training group.
- For all walking outcomes, 57% to 80% of participants in the high-intensity groups had important clinical gains, while only 9% to 31% of participants did so following low-intensity training.
- High-intensity variable training also resulted in improved dynamic balance while walking and improved balance confidence.

Hornby noted that no serious adverse events occurred during the training sessions, suggesting [stroke survivors](#) can be pushed to higher intensity walking with more variable tasks during rehabilitation.

"Rehabilitation that allows walking practice without challenging the nervous system doesn't do enough to make a statistical or clinically significant difference in a patient's recovery after a stroke," Hornby said. "We found that when [stroke patients](#) are pushed harder, they see greater changes in less time, which translates into more efficient rehabilitation services and improved mobility."

Ultimately, their goal is to incorporate high-intensity variable step training into regular clinical

[rehabilitation](#) protocols.

The study was small compared to larger, multicenter clinical trials. Hornby said the next step would be to test high-intensity, variable step [training](#) in larger patient populations in a large, multicenter clinical trial.

More information: T. George Hornby et al, Contributions of Stepping Intensity and Variability to Mobility in Individuals Poststroke, *Stroke* (2019).
[DOI: 10.1161/STROKEAHA.119.026254](https://doi.org/10.1161/STROKEAHA.119.026254)

Provided by American Heart Association

APA citation: High-intensity step training boosts stroke survivors' walking skills (2019, August 22) retrieved 28 August 2022 from <https://medicalxpress.com/news/2019-08-high-intensity-boosts-survivors-skills.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.