

NIH study suggests higher air pollution exposure during second pregnancy may increase preterm birth risk

13 September 2019



Credit: CC0 Public Domain

Pregnant women who are exposed to higher air pollution levels during their second pregnancy, compared to their first one, may be at greater risk of preterm birth, according to researchers at the National Institutes of Health. Their study appears in the *International Journal of Environmental Research and Public Health*.

Preterm birth, or the birth of a baby before 37 weeks, is one of the [leading causes](#) of infant mortality in the United States, according to the Centers for Disease Control and Prevention. Although previous studies have found an association between [air pollution exposure](#) and preterm birth risk, the authors believe their study is the first to link this risk to changes in exposure levels between a first and second pregnancy.

"What surprised us was that among low-risk women, including women who had not delivered preterm before, the risk during the second pregnancy increased significantly when air pollution stayed high or increased," said Pauline

Mendola, Ph.D., the study's lead author and a senior investigator in the Epidemiology Branch at the Eunice Kennedy Shriver National Institute of Child Health and Human Development.

Researchers used data from the NICHD Consecutive Pregnancy Study to examine the risk of preterm birth. They matched [electronic medical records](#) of more than 50,000 women who gave birth in 20 Utah hospitals between 2002 and 2010 to data derived from Community Multiscale Air Quality Models, modified based on a model by the Environmental Protection Agency, which estimate pollution concentrations.

Researchers examined exposure to sulfur dioxide, ozone, [nitrogen oxides](#), nitrogen dioxide, carbon monoxide and particles. For nearly all pollutants, exposure was more likely to decrease over time, but 7 to 12 percent of women in the study experienced a higher exposure to air pollution during their second pregnancy. The highest risks were with increasing exposure to carbon monoxide (51 percent) and nitrogen dioxide (45 percent), typically from emissions from motor vehicles and power plants; ozone (48 percent), a secondary pollutant created by combustion products and sunlight; and [sulfur dioxide](#) (41 percent), mainly from the burning of fossil fuels that contain sulfur, such as coal or diesel fuel.

More research is needed to confirm this association, but improvements in air quality may help mitigate [preterm birth](#) risk among [pregnant women](#), Dr. Mendola said.

More information: Pauline Mendola et al. Air Pollution and Preterm Birth: Do Air Pollution Changes over Time Influence Risk in Consecutive Pregnancies among Low-Risk Women?, *International Journal of Environmental Research*

and Public Health (2019). DOI:
[10.3390/ijerph16183365](https://doi.org/10.3390/ijerph16183365)

Provided by National Institutes of Health
APA citation: NIH study suggests higher air pollution exposure during second pregnancy may increase preterm birth risk (2019, September 13) retrieved 13 November 2022 from
<https://medicalxpress.com/news/2019-09-nih-higher-air-pollution-exposure.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.