

Incidence of Hodgkin lymphoma up in least deprived areas

September 26 2019



Individuals living in the most affluent areas of the United Kingdom seem

to have an increased incidence of Hodgkin lymphoma (HL), according to a study published in the August issue of *BMJ Open*.

Meena Rafiq, M.B.Ch.B., from University College London, and colleagues analyzed data on 10 million individuals in the United Kingdom from 1992 to 2016 to identify trends and high-risk populations.

The researchers identified 2,402 new cases of HL during 78,569,436 person-years. Significant variation was seen in HL incidence by [deprivation](#) group. Compared with those individuals living in the most deprived areas, those in the most affluent areas had a significantly higher HL incidence (incidence rate ratio, 1.60), with strong evidence of a linear trend toward increasing HL incidence with decreasing deprivation. Across the United Kingdom, there was significant regional variation in HL incidence, which persisted after adjustment for age, sex, and deprivation (incidence rate ratio, 0.80 to 1.42).

"The findings are consistent with the hypothesis that an affluent childhood environment may predispose to development of immune-related conditions, possibly through fewer immune challenges interfering with the maturation of the immune system," the authors write.

More information: [Abstract/Full Text](#)

Copyright © 2019 [HealthDay](#). All rights reserved.

Citation: Incidence of Hodgkin lymphoma up in least deprived areas (2019, September 26) retrieved 10 May 2023 from <https://medicalxpress.com/news/2019-09-incidence-hodgkin-lymphoma-deprived-areas.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.