

Weight-loss surgery cuts risk of birth defects

October 15 2019



Martin Neovius, professor in the Department of Medicine in Solna at Karolinska Institutet in Sweden. Credit: Kate Gabor

Children born to women who underwent gastric bypass surgery before becoming pregnant had a lower risk of major birth defects than children

born to women who had severe obesity at the start of their pregnancy. That's according to a matched cohort study by researchers at Karolinska Institutet and Örebro University published in the scientific journal *JAMA*. The findings indicate that weight-loss and improved blood sugar control could reduce the risk of major birth defects and ought to lessen longstanding concerns that weight-loss surgery could increase this risk.

Obesity and poor [blood sugar control](#) have in previous studies been linked to an increased risk of health complications for both mothers and their infants. Other studies have shown that [weight-loss surgery](#) may increase the risk of various nutrient deficiencies, including iron and folate which are important for fetal development. There have been concerns dating back to the 1980s that [bariatric surgery](#) could increase the risk of major birth defects. In recent years, weight-loss surgery has increased significantly, and about 1.5 percent of all babies born in Sweden today are delivered by mothers who have had bariatric surgery.

The current study shows, however, that the risk of major birth defects was about 30 percent lower in children of mothers who had [gastric bypass surgery](#) than in children of severely obese mothers. The risk of major defects was 3.4 percent in children born to [women](#) who had had gastric bypass surgery, which is in line with the risk of major defects in children born to normal weight women (3.5 percent). For women who at their first prenatal checkup had a body-mass index comparable to that of the gastric bypass patients' pre-surgery weight, the risk of major birth defects was 4.9 percent.

"This study shows that weight-loss and improved blood sugar control in the mother can actually result in a lower risk of birth defects in the child," says Martin Neovius, professor and researcher at the Department of Medicine in Solna and one of the study's main authors. "It should help reduce fears that bariatric surgery increases the risk of birth defects in the event of future pregnancy, assuming that surgery patients take their

recommended [nutritional supplements](#)."

Lost 40 kilos

To compare the two groups, the researchers examined data on more than 33,000 births in Sweden between 2007 and 2014. Of these, 2,921 were children born to mothers who underwent a gastric bypass, and 30,753 were children born to women who weighed about the same as the gastric bypass women did before their surgeries. The women in the surgery group lost an average of 40 kilos and weighed around 82 kilos at their first prenatal checkup. Their use of diabetes medication decreased to 1.5 percent from 9.7 percent.

"Bariatric surgery has many [positive effects](#) on pregnancy," says Olof Stephansson, obstetrician and senior researcher at the Department of Medicine in Solna, and one of the study's main authors. "Still, it is important that these women receive special antenatal care, including additional ultrasounds to monitor fetal growth and detailed nutritional counseling that include administration of supplements necessary after weight-loss [surgery](#). Antenatal care providers should also check for nutrition deficiencies in addition to iron, such as folic acid, calcium and vitamin B12."

More information: "Association of Maternal Gastric Bypass Surgery with Offspring Birth Defects," *JAMA* (2019). [DOI: 10.1001/jama.2019.12925](#)

Provided by Karolinska Institutet

Citation: Weight-loss surgery cuts risk of birth defects (2019, October 15) retrieved 18 July 2023 from <https://medicalxpress.com/news/2019-10-weight-loss-surgery-birth-defects.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.