


Unhealthy habits can start young: Infants, toddlers, and added sugars

November 14 2019



Top eight sources of added sugars in the diets of US infants and toddlers aged 6 to 23 months, National Health and Nutrition Examination Survey, 2011-2016. Credit: CDC/NCHS, National Health and Nutrition Examination Survey, 2011-2016

A new study in the *Journal of the Academy of Nutrition and Dietetics* found that nearly two-thirds of infants (61 percent) and almost all toddlers (98 percent) consumed added sugars in their average daily diets,

primarily in the form of flavored yogurts (infants) and fruit drinks (toddlers). Infants were 6-11 months, and toddlers were 12-23 months.

The analysis documented some good news in the decline over the study period (2005-6 and 2015-16) in the percentage of infants and toddlers whose daily diets include added sugars, as well as the amounts they consumed. Yet the widespread intake points to a serious and persistent problem: the [early development](#) of eating patterns associated with negative health conditions.

"Our study, which is the first to look at trends in added sugars consumption by infants and toddlers, documents that most infants and toddlers consume added sugars. This has important public health implications since previous research has shown that eating patterns established early in life shape later eating patterns," explained lead investigator Kirsten A. Herrick, Ph.D., MSc, Division of Health and Nutrition Examination Surveys, National Center for Health Statistics, Centers for Disease Control and Prevention, Hyattsville, MD, USA. She cited an earlier study that found that 6-year-olds who had consumed any [sugar](#)-sweetened beverage (SSB) before the age of one were more than twice as likely to consume an SSB at least once a day compared to 6-year-olds who had not consumed any before the age of one.

Dr. Herrick noted, "Previous research into the diets of children over two years old associated sugar consumption with the development of cavities, asthma, obesity, elevated blood pressure and altered lipid profiles."

Health organizations in the United States promulgate guidelines that recommend limiting sugar intake to 9 tsp or less for adult men, and 6 tsp or less for adult women and children between 2 and 19. With no comparable research available for infants and toddlers prior to this study, only one organization, the American Heart Association, provided any guidance for children under age 2. "Our study's findings about infant and

[toddler](#) diets should raise awareness among health organizations and practitioners and inform future guidelines and recommendations," she added.

The investigators analyzed data for 1,211 infants and toddlers (6-23 months) from the National Health and Nutrition Examination Survey (NHANES) 2011-2016, a nationally representative study conducted by the US Centers for Disease Control and Prevention. They used the Food Patterns Equivalents Database and the United States Department of Agriculture's What We Eat In America's list to categorize foods. Sugars contained in [breast milk](#) and formula were not included in the consumption estimates.

The results showed that infants consumed about 1 teaspoon (tsp) of added sugars daily (equivalent to about 2 percent of their daily caloric intake), while toddlers consumed about 6 tsp of sugars (about 8 percent of their daily caloric intake). No differences were detected in added sugars consumption by sex, family income level, and head-of-household, but there were some distinctions by race/Hispanic origin: non-Hispanic Asian toddlers consumed the fewest added sugars (3.7 tsp) and non-Hispanic black toddlers consumed the most added sugars (8.2 tsp). The top food sources of added sugars for infants included yogurt, baby snacks and sweets, and sweet bakery products. For toddlers, the top sources included fruit drinks, sweet baked products, and sugar and candy.

According to Dr. Herrick, parents should be mindful of added sugars levels in the foods chosen when weaning their [infants](#). "The transition from a milk-based diet (breast milk and formula) to table foods has an impact on nutrition, taste preference, and eating patterns. More work is needed to understand this critical period." She recommends discussing which solid foods to introduce during weaning with a child's healthcare provider and pointed to the Nutrition Facts label as another resource to

support informed decisions. While the federal requirement to include added sugars content of a food or beverages on the Nutrition Facts label is not mandatory until January 2020, many labels already include this information. For more information about infant and toddler nutrition, visit <http://www.cdc.gov/nutrition>.

In September 2019, the Academy of Nutrition and Dietetics joined the American Heart Association, the American Academy of Pediatric Dentistry, and the American Academy of Pediatrics to recommend breast milk, infant formula, water, and plain milk as part of a new set of comprehensive beverage recommendations for children, outlined by age (birth through age 5). They caution against beverages that are sources of added sugars in young children's diets, including flavored milks (e.g., chocolate, strawberry) and sugar- and low-calorie sweetened beverages, in addition to a wide variety of beverages that are on the market and targeted to children that provide no unique nutritional value.

More information: Added Sugars Intake among US Infants and Toddlers, *Journal of the Academy of Nutrition and Dietetics* (2019). [DOI: 10.1016/j.jand.2019.09.007](https://doi.org/10.1016/j.jand.2019.09.007)

Provided by Elsevier

Citation: Unhealthy habits can start young: Infants, toddlers, and added sugars (2019, November 14) retrieved 4 February 2023 from <https://medicalxpress.com/news/2019-11-unhealthy-habits-young-infants-toddlers.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.