


Implant survival poor for converted 'partial' knee replacements

3 December 2019



[survival](#) when compared with that following primary TKA," the authors write.

More information: [Abstract/Full Text](#)
(subscription or payment may be required)

Copyright © 2019 [HealthDay](#). All rights reserved.

(HealthDay)—Total knee arthroplasty (TKA) converted from medial unicompartamental (or "partial") knee arthroplasty (UKA) has a risk for revision that is threefold higher than that for primary TKA, according to a study published online Nov. 20 in *The Journal of Bone & Joint Surgery*.

Anders El-Galaly, M.D., from Aalborg University Hospital in Denmark, and colleagues used data from the Danish Knee Arthroplasty Registry (1997 to 2017) to assess survival of 1,012 TKAs converted from UKAs versus 73,819 primary TKAs and 2,572 revision TKAs.

The researchers found that the survival of TKAs converted from UKAs was comparable with that of revision TKAs ($P = 0.42$) and significantly inferior to the survival of primary TKAs (P

"On the basis of this study, we believe that careful consideration is necessary before using medial UKA as treatment for [knee](#) osteoarthritis, as a potential conversion to a TKA decreased implant

APA citation: Implant survival poor for converted 'partial' knee replacements (2019, December 3)
retrieved 12 October 2022 from <https://medicalxpress.com/news/2019-12-implant-survival-poor-partial-knee.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.