

How do we detect if coronavirus is spreading in the community?

4 March 2020, by Joanne MacDonald



Credit: Shutterstock

The daily [number of new coronavirus \(SARS-CoV-2\)](#) cases is now [nine times higher outside China](#) than in the country where the disease was first detected.

In Australia, [reports this week of local transmission](#) of the coronavirus, which causes the [disease](#) now called COVID-19, are a turning point in our disease management.

Our disease [surveillance](#) systems are well placed to keep abreast of COVID-19 and provide some reassurance that transmission is unlikely to go undetected in the community. But there's still more we could do.

Why is the coronavirus hard to detect?

Detecting all people with COVID-19 is a global problem. There are too many existing respiratory infections, such as colds and flus, in any country to be able to test everyone with coronavirus symptoms.

Each of us gets around two to three upper respiratory tract infections a year. Globally, that amounts to around [around 18.75 billion infections a year](#).

There are not enough testing kits readily available to test people at this scale.

So who is tested?

In Australia, people are [currently tested](#) for the coronavirus if they've travelled from or through a country considered to pose a risk of transmission in the 14 days before getting sick, or if they have a link to a known case.

This testing criteria has changed as the outbreak progressed, and will continue to do so.

How else do we track possible cases?

Apart from directly testing people suspected of having COVID-19, Australia has a [surveillance plan](#) to detect coronavirus in people or populations who don't know they're infected.

Australia's [emergency response plan for mitigating COVID-19](#) says we will use surveillance networks set up for the [influenza pandemic emergency reponse plan](#) and some of this is already occurring.

The [Australian Sentinel Practices Research Network](#) (ASPREN) is a network of GPs who log the number of patients they see in total, compared to the number of patients they see with influenza-like illness. These GPs collect samples from a small subset of those patients to see if COVID-19 is circulating. Samples are then sent to [SA Pathology](#) for testing.

Another surveillance network that may be activated is FluCAN. This reports on the number of hospitalisations due to a disease, usually influenza, as well as [clinical data](#) from the cases. The information helps public health experts get a better picture of how severe the disease is, and the symptoms.

But while these systems can monitor disease levels

in those sick enough to seek medical care, they don't give us an indication of the amount of milder disease that might be circulating in our communities.

This is where an online surveillance system called [FluTracking](#) can help. Anyone in the community can join and answer two simple questions each week about whether they have a fever and/or a cough.

The system provides information on how much influenza-like illness is circulating in the community. If we're seeing more than usual, it might signify a community outbreak.

FluTracking was activated for COVID-19 surveillance [last month](#).

What else could we be doing?

There are questions, however, about how early in an outbreak the surveillance systems will detect cases.

Will they detect community transmission when an outbreak reaches ten cases? Or will it take hundreds or even thousands of cases to trigger a warning through the network?

Innovations from other countries suggest there are additional measures Australia could adopt to improve our surveillance networks.

Testing at home or on the road

London adopted a system at the end of January to test mild cases of disease in the [patients' homes](#). This helps with self-isolation and reducing disease spread.

Meanwhile, Edinburgh has opened a [drive-through testing clinic](#) to reduce the chances of viral spreading.

Australia's emergency response plans include provisions to mobilise flu clinics to help keep patients from overwhelming [emergency services](#). Drive-through services could be an excellent addition to these existing plans.

Rapid testing

One of the biggest concerns with current testing is the time it takes to ship a sample to a laboratory for testing. This can result in a delay of one to two days before getting the results. During peak epidemic times, testing can't cope with demand.

Researchers in China, however, have reportedly developed a [rapid coronavirus test](#) that can detect the virus using a fingerprick of blood in 15 minutes. The [test](#) detects if the body has mounted an immune response (Ig M antibodies) to the virus.

While the data is not yet published, the researchers reported success [from the 600 samples they tested](#).

Rapid diagnostic tests are typically cheap to manufacture, can be mass produced, and can be easily used by health workers outside a laboratory.

Reporting on negative test results and surveillance systems

Positive COVID-19 cases from all the surveillance sources are [reported as they occur](#). But while influenza surveillance reports include the number of negative tests results, it's unclear whether COVID-19 surveillance reports will do the same.

Reporting on negative tests results could help ease community concern that coughs and sneezes people see on the train or in their office are unlikely to be due to the undetected spread of COVID-19.

Local health services should also provide regular updates on the types of COVID-19 surveillance actively being performed, and in which communities. This information can help reduce anxiety levels and assure communities that spread of [coronavirus](#) can be effectively monitored.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

APA citation: How do we detect if coronavirus is spreading in the community? (2020, March 4) retrieved 3 December 2022 from <https://medicalxpress.com/news/2020-03-coronavirus.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.