


# Stroke: When the system fails for the second time

24 March 2020



The second lesion, in which large parts of the left hemisphere are not working anymore (A, dark grey), increased the contribution of the right brain. The individual impairment predicted the activation on the right side (A, yellow). The stronger the fibre connection between the sister areas (B, red) on the right side, the less the patient was affected by the interruption on the left. Credit: MPI CBS

After a stroke, there is an increased risk of suffering a second one. If areas in the left hemisphere were affected during the first attack, language is often impaired. In order to maintain this capability, the brain usually briefly drives up the counterparts on the right side. But what happens after a second attack? The Max Planck Institute for Human Cognitive and Brain Sciences has found an answer using virtual lesions.

The [brain](#) is much more malleable than once thought. Even after stroke or [brain injury](#), the brain often succeeds finding a new balance between the failed regions and the functions they serve. Commonly, neighboring regions are activated as well as homologues on the other side of the brain side. During language processing, the homologues of the left-dominant language areas are usually less active and are kept in check by the dominant half—until the emergency case occurs.

Until now, it was unclear whether these mechanisms also apply in the event of a second attack. Does the brain retain its capacity to adapt? This is important, as up to 15 percent of those

affected will have a second stroke. In addition, there was disagreement about whether an activated right brain is generally good for healing. Some studies suggest that involvement of the right [hemisphere](#) helps recovery, at least in the short term. Others found, however, that a loss of language areas in the left half can literally inhibit the right half. In that case, the contribution of the right hemisphere has nothing to do with language and can cause confusion. The brain gets out of step. Further, studies had also found that the patients are better off if the overactive half is restrained by inhibitory magnetic stimulation, in which case the activity shifts back to the [left hemisphere](#) and it wins the upper hand again.

Scientists at the Max Planck Institute for Human Cognitive and Brain Sciences (MPI CBS) in Leipzig, Germany, have now found that the brain areas on the right side also become more active when there is a second injury in the left language areas. "In the recovered brain, the right side's contribution was still little after the first impairment. After the second event, in which large parts of the left hemisphere are not working anymore, its role becomes much more important," explains Gesa Hartwigsen, research group leader at the MPI CBS and first author of the article, which has been published in the high-ranking, open access journal *eLife*. "The second lesion increased the contribution of the right brain," Hartwigsen said.

The scientists examined these relationships using 12 patients in whom the regions for processing properties of sound in the left hemisphere were injured. The incident had happened to them at least six months prior. Their brain had the opportunity to regenerate and adapt to the new situation. The researchers simulated the second disruption using so-called transcranial magnetic stimulation, which can be used to briefly halt certain areas of the brain through electrical stimuli. It can be used to simulate how the brain would react if certain areas actually fail due to a stroke or other events—and how this

affects the ability to recognize sounds, for example. To do this, Hartwigsen and her team used a simple decision task. The participants heard the word "cat" and had to decide whether it consisted of one or two syllables. The individual impairment predicted the activation on the right side. The researchers also found that the stronger the fiber connection between the sister areas on the right side, the less the patient was affected by the interruption on the left.

"These results show that after large-scale disturbances in which large parts of the left hemisphere no longer function as they should, the right hemisphere probably plays a beneficial role. Often, there is a lot of tissue in the left half of the brain that only works to a limited extent and needs support from the right side. "Other studies show that recovery is helped when the activated right side later down-regulates itself and thus contributes to normalization on the left side," said Hartwigsen. On the other hand, if the right half remains permanently up-regulated, healing is delayed.

Findings on how the damaged brain adapts to repeated injury could help to improve the therapy of stroke patients in the long term. "This may make it possible, at some point, to assess whether it would be more helpful to regulate specific areas up or down," says Hartwigsen, confidently.

**More information:** Gesa Hartwigsen et al, Short-term modulation of the lesioned language network, *eLife* (2020). [DOI: 10.7554/eLife.54277](https://doi.org/10.7554/eLife.54277)

Provided by Max Planck Society

APA citation: Stroke: When the system fails for the second time (2020, March 24) retrieved 30 May 2022 from <https://medicalxpress.com/news/2020-03-stroke-when-the-system-fails.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*