

Shorter radiotherapy treatment for bowel cancer patients during COVID-19

6 April 2020



Professor David Sebag-Montefiore. Credit: University of Leeds

An international panel of cancer experts has recommended a one-week course of radiotherapy and delaying surgery as the best way to treat patients with bowel cancer during the COVID-19 pandemic.

The short course of treatment involves higher-intensity radiation rather than five weeks of [radiotherapy](#) coupled with chemotherapy. Surgery, which normally happens one to two weeks after radiotherapy, can be safely delayed by up to 12 weeks, say the [expert](#) panel.

This approach, based on the latest research evidence, will maintain the best chance of successfully treating the disease while at the same time reducing the side effects of treatment and the risks of COVID-19 infection.

People with [bowel cancer](#) are more susceptible to severe complications from COVID-19 because their immune system is weakened. Shorter-course radiotherapy avoids the need for chemotherapy, which further suppresses the immune system. It

also means significantly fewer hospital appointments, allowing patients to maintain social distancing rules.

David Sebag-Montefiore, Professor of Clinical Oncology at the University of Leeds and Honorary Clinical Oncologist with the Leeds Teaching Hospitals NHS Trust, who led the [expert panel](#), said: "The COVID-19 pandemic is a global emergency and we needed to work very quickly to identify changes that would benefit patients. Our recommendations were published 20 days after our first meeting.



Professor David Sebag-Montefiore. Credit: University of Leeds

"This process normally takes many months, if not years."

Writing in the journal *Radiotherapy and Oncology*, the panel, made up of cancer experts from across Europe, say it is also possible that hospitals may struggle to offer the current treatment approaches

as COVID-19 impacts on hospital staffing levels.

The 15-strong panel comprised researchers who led the defining studies. Their research shows that surgery can be safely delayed by 12 weeks. The chances of successful treatment are maintained and post-operative side effects are reduced. This allows surgery to be scheduled after the peak of the pandemic.

The recommendation to use shorter-course radiotherapy follows a major study, funded by the UK Medical Research Council and led by Professor Sebag-Montefiore, which demonstrated the benefit of the one-week course of radiotherapy.

Professor Sebag-Montefiore added: "Our guidelines will result in a very substantial change in treatment across the globe. During the COVID-19 pandemic, our patients will benefit from the use of an effective, shorter and safer radiotherapy treatment."

More information: The expert panel's recommendations can be found here:

www.thegreenjournal.com/article/10.1002/1473-2165.10140

Provided by University of Leeds

APA citation: Shorter radiotherapy treatment for bowel cancer patients during COVID-19 (2020, April 6) retrieved 29 April 2021 from

<https://medicalxpress.com/news/2020-04-shorter-radiotherapy-treatment-bowel-cancer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.