

Study links stress of a marathon on the heart to the intensity at which it is run

28 July 2020



Credit: CC0 Public Domain

The number of recreational/non-elite athletes participating in marathons is gradually increasing every year. The great physical effort experienced by their body in this type of competition triggers an acute release of biomarkers of heart damage (cardiac troponins), making them rise above clinically significant values; that is, reaching levels above the 99th percentile of a healthy reference population.

The *European Journal of Applied Physiology* has published a study led by Dr. Ignacio Martínez, Doctor in Physical Activity and Sport Sciences and member of the team of the Sports Health Unit of the Vithas Valencia 9 de Octubre hospital, which shows that cardiac stress caused by a marathon is proportional to the relative intensity at which it is run. This implies that the athletes who are able to perform this type of competition at a higher percentage with respect to their maximum heart rate, show higher concentrations of heart damage markers in their blood at the end of the race.

Furthermore, research shows that, on the contrary, neither echocardiographic variables nor

cardiovascular fitness show any relationship with the release of troponin, a marker of heart damage. "Compared to similar studies carried out in this field, our work reveals, for the first time, that 24 hours after finishing the marathon, those runners who still show a concentration of troponin in their blood above the reference values are those who ran the marathon at a higher relative speed," assures Dr. Ignacio Martínez.

The study was carried out at the 2016 Valencia Marathon on a sample of 98 runners, who were subjected to a stress test and an echocardiogram before the race. Likewise, [blood samples](#) were obtained from each of the athletes before, immediately after and at 24, 48, 96, 144 and 192 hours after the race. The results showed a significant increase in heart damage markers at the end of the [marathon](#) and a normalization of these markers linked to the relative speed of the race.

More information: Ignacio Martínez-Navarro et al. Immediate and 24-h post-marathon cardiac troponin T is associated with relative exercise intensity, *European Journal of Applied Physiology* (2020). [DOI: 10.1007/s00421-020-04403-8](https://doi.org/10.1007/s00421-020-04403-8)

Provided by Asociacion RUVID

APA citation: Study links stress of a marathon on the heart to the intensity at which it is run (2020, July 28) retrieved 27 July 2022 from <https://medicalxpress.com/news/2020-07-links-stress-marathon-heart-intensity.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.