


# Study tracks chlamydia trachomatis infection in Sweden

22 February 2021



Chlamydia trachomatis inclusion bodies (brown) in a McCoy cell culture. Credit: public domain

Chlamydia trachomatis infection is the most common bacterial sexually transmitted infection (STI), with about 127 million new cases estimated annually worldwide. Since it is an asymptomatic infection, individuals may carry it for a long time and unknowingly transmit the infection to others.

Untreated [chlamydia](#) may lead to serious sequelae of the reproductive tract which is why early detection of infected individuals via testing, treatment, and partner notification is vital and may prevent further transmission of infection.

In her thesis, Inga Veli?ko aimed to extend our understanding of chlamydia epidemiology at an individual and population-based level. The thesis is based on the data from a cohort study in the urban STI clinic, as well as chlamydia cases and tests reported to the national infectious diseases register SmiNet-2 at the Public Health Agency of Sweden.

The risk factors independently associated with chlamydia diagnosis identified in Inga's work (being 20-24 years old, having six or more [sexual](#)

[partners](#) during the previous 12 months, using alcohol before sex, reporting all type of sexual activities during the last sexual contact, and testing due to partner notification) are in line with previous studies. Her results also suggest that individuals in the high-risk sexual behavior latent classes are more likely to [test](#) repeatedly for chlamydia, suggesting absorption of public health messages. Inga also found that testing volumes were driving chlamydia notification rates during time of 1992-2004, but not later during 2009 to 2018. This suggested that notification rates during the latter period were less biased and may suggest a true decrease in chlamydia incidence. In the last thesis study, Inga found a decrease in prevalence which could explain the observed decrease in chlamydia [notification](#) rates during 2009-2018.

Inga concludes that prevention work should be continued to reach asymptomatic individuals by testing and prevention messages as her results suggest that these individuals practice unsafe sex. Moreover, her results suggested that observed decrease in chlamydia rates could be due to an actual decrease in chlamydia prevalence.

On March 12, Inga will defend her thesis "Chlamydia trachomatis [infection](#) in Sweden: time trends, [risk factors](#), and prevalence."

**More information:** Chlamydia trachomatis infection in Sweden: time trends, risk factors, and prevalence. [openarchive.ki.se/xmlui/handle/10616/47497](https://openarchive.ki.se/xmlui/handle/10616/47497)

Provided by Karolinska Institutet

APA citation: Study tracks chlamydia trachomatis infection in Sweden (2021, February 22) retrieved 30 April 2021 from <https://medicalxpress.com/news/2021-02-tracks-chlamydia-trachomatis-infection-sweden.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*