


# Diphtheria immunization in WHO European region a success, but still some sporadic cases

February 24 2022

Number of reported diphtheria cases<sup>a</sup> (n=451) and DTP1 and DTP3 coverage in the WHO European Region, 2010–2019



DTP1: first dose of diphtheria, tetanus and pertussis vaccine; DTP3: third dose of diphtheria, tetanus and pertussis vaccine; WHO: World Health Organization.

<sup>a</sup> The number of countries that submitted reports (including zero reporting) on diphtheria cases are shown in parentheses below the year.

For 2010–19, the number of countries that submitted reports on diphtheria cases in the WHO European Region (including zero reporting) ranged from 38 in 2014 to 53 in 2018. During this 10-year period, there were 451 cases of diphtheria reported in the Region. Credit: Eurosurveillance

In their article published in *Eurosurveillance* today, Muscat et al. present diphtheria data on the number of notified cases between 2010 and 2019 from the WHO/United Nations International Children's Emergency Fund (UNICEF) Joint Reporting Forms and analyze it by age and vaccination status.

Diphtheria re-emerged in the WHO European Region in the 1990s in the newly independent states of the former Soviet Union. Today, diphtheria is considered uncommon following successful mass immunization with only sporadic reports of cases in the WHO European Region, however, some of them fatal. Immunization with the available diphtheria vaccines is the only effective method of preventing this acute bacterial disease which was previously a major cause of death among children.

## **Good vaccination coverage explains relatively few cases**

Worldwide, over 87,500 diphtheria cases have been reported between 2010 and 2019. During the same 10-year period, the WHO European Region recorded only 451 diphtheria cases. In 2019, 52 cases were notified by 11 of 48 countries that submitted data. Although relatively few, the cases observed in the European Region are of concern, and may be also partly attributed to suboptimal surveillance systems and inadequate or lack of specialized laboratory diagnostic capacity.

Coverage with the third vaccine dose (DTP3) was relatively stable at a high level in the period 2010-2019 across the Region. For 2019, it was at 95% with 32 out of the 53 countries of the Region reporting more than 95% coverage and only six countries reporting coverage below 90%. For that same year, 26 countries achieved a high ( $\geq 90\%$ ) DTP3 coverage in more than 90% of their districts. Twenty-eight countries reported none of their districts below 80% coverage with the third dose (DTP3). Seven countries had 1–5% of their districts with less than 80% DTP3

coverage.

## **Adults aged 30 and older most affected**

For the year 2019, where more detailed data were available, the age of reported cases ranged from less than a year old to older than 30 years of age. Most diphtheria were reported in adults aged 30 years and older. This age distribution shift with cases mainly among adolescents and adults concurs with recent review findings of diphtheria epidemiology.

## **Ensuring adequate laboratory capacity to diagnose toxigenic diphtheria is crucial**

The authors conclude that laboratory testing of all suspected cases should be conducted for case confirmation. An adequate surveillance of diphtheria requires that laboratories are equipped with the appropriate materials and that all isolates of potentially toxigenic corynebacteria should ideally be submitted to a reference/specialist laboratory for confirmation of identification and toxigenicity testing. A revised World Health Organization manual for laboratory diagnosis of diphtheria and related infections has recently been published to assist laboratory workers in the correct procedures to diagnose diphtheria cases and to guide clinicians in [treatment options](#).

The authors conclude that "long-standing high DTP3 coverage at Regional level probably explains the relatively few diphtheria cases reported in the Region. Suboptimal surveillance systems and inadequate laboratory diagnostic capacity may also be contributing factors. Still, the observed cases are of concern. Attaining high DTP3 coverage in all districts and implementing recommended booster doses are necessary to control [diphtheria](#) and prevent outbreaks."

**More information:** Mark Muscat et al, Diphtheria in the WHO

European Region, 2010 to 2019, *Eurosurveillance* (2022). [DOI: 10.2807/1560-7917.ES.2022.27.8.2100058](https://doi.org/10.2807/1560-7917.ES.2022.27.8.2100058)

Provided by European Centre for Disease Prevention and Control (ECDC)

Citation: Diphtheria immunization in WHO European region a success, but still some sporadic cases (2022, February 24) retrieved 18 July 2023 from <https://medicalxpress.com/news/2022-02-diphtheria-immunization-european-region-success.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.